





COVID-19 Vaccination Schedule for people who are **NOT** Moderately or Severely Immunocompromised


People ages 6 months through 4 years




-OR-




People age 5 years




-OR-




People ages 6 through 11 years



People ages 12 years and older



People ages 18 years and older who previously received Janssen primary series dose‡



*For people who previously received a monovalent booster dose(s), the bivalent booster dose is administered at least 2 months after the last monovalent booster dose.


† A monovalent Novavax booster dose maybe used in limited situations in people 18 years and older who completed a primary series using any COVID-19 vaccine, have not received any previous booster dose(s), and are unable or unwilling to receive an mRNA vaccine. The monovalent Novavax booster dose is administered at least 6 months after completion of a primary series.

‡Janssen COVID-19 dose should only be used in certain limited situations. See: <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a>




COVID-19 Vaccination Schedule for people who **ARE** Moderately or Severely Immunocompromised


People ages 6 months through 4 years




People age 5 years




People ages 6 through 11 years



People ages 12 years and older




People ages 18 years and older who previously received Janssen primary series dose[‡]



Monoclonal antibodies (EVUSHELD™) for COVID-19 pre-exposure prophylaxis

People ages 12 years and older (must weigh at least 40kg)



*For people who previously received a monovalent booster dose(s), the bivalent booster dose is administered at least 2 months after the last monovalent booster dose.

†A monovalent Novavax booster dose maybe used in limited situations in people 18 years and older who completed a primary series using any COVID-19 vaccine, have not received any previous booster dose(s), and are unable or unwilling to receive an mRNA vaccine. The monovalent Novavax booster dose is administered at least 6 months after completion of a primary series.

‡Janssen COVID-19 dose should only be used in certain limited situations. See: <https://www.cdc.gov/vaccines/covid-19/clinical-considerations/interim-considerations-us-appendix.html#appendix-a>